



What data are needed for blood collectors to provide **cold-stored** whole blood?


Rebecca Cardigan PhD, FRCPath

Head of Component Development/PI, NHS Blood & Transplant, England

Affiliated Lecturer, Dept Haematology, University of Cambridge

Caring Expert Quality

Conflicts of interest


- I am a member of Terumo BCT's Advisory Board, who make filters for LD whole blood
 - I work for a national blood provider!
- 
- A thick, solid blue line that curves across the bottom of the slide, starting from the left edge, dipping down, and then rising towards the right edge.

Content


- What is the clinical need/benefit v current standard of care?
- Donor selection
- Manufacture and storage of the product
- Cost and logistics
- What are the outstanding research questions?

These considerations differ depending on current standard of care, nature of blood supplier & local societal considerations in relation to risk

Pre-hospital treatment of major haemorrhage has evolved



Pre-hospital protocols for resuscitation vary



clear fluids only



RBC



RBC + thawed/dried plasma



RBC + thawed/dried plasma + cold stored platelets



Whole blood


Perceived benefits of whole blood transfusion



Balanced resuscitation
1:1:1



Improved
efficacy/reduced
transfusion
requirements



Simpler/reduced
time to administer

Can we reduce the mortality of patients from traumatic major haemorrhage?

Evidence that WB is superior to component therapy is lacking

> Emerg Med J. 2020 Jun;37(6):370-378. doi: 10.1136/emmermed-2019-209040. Epub 2020 May 6.

Whole blood transfusion versus component therapy in adult trauma patients with acute major haemorrhage

Pascale Avery^{1 2}, Sarah Morton³, Harriet Tucker^{2 4 5}, Laura Green^{2 6 7}, Anne Weaver^{2 8 9}, Ross Davenport^{2 10}



Review

Assessing the risks of haemolysis as an adverse reaction following the transfusion of ABO incompatible plasma-containing components - A scoping review

Josephine McCullagh^{a,b,1,*}, Rebecca Cardigan^{c,g,1}, Susan J. Brunskill^f, Tom Bullock^d, Carolyn Doree^f, Lise Estcourt^f, Sian Huish^c, Josie Sandercock^f, Laura Green^{a,c,e}

^a Barts Health NHS Trust, London, United Kingdom

^b Bolton NHS Foundation Trust, Greater Manchester, United Kingdom

^c Component Development Laboratory, NHS Blood and Transplant, Cambridge, United Kingdom

^d Red Cell Immunohaematology NHS Blood and Transplant, United Kingdom

^e Blizard Institute, Queen Mary University of London, London, United Kingdom

^f Systematic Review Initiative, NHS Blood and Transplant, Oxford, United Kingdom

^g Department of Haematology, University of Cambridge, United Kingdom

SYSTEMATIC REVIEW

Outcome measures used in clinical research evaluating prehospital blood component transfusion in traumatically injured bleeding patients: A systematic review

Tucker, Harriet FRCEM FIMC RCSEd; Avery, Pascale BMBS, MSc; Brohi, Karim FRCS, FRCA; Davenport, Ross FRCS, PhD; Griggs, Joanne MSc; Weaver, Anne FCEM; Green, Laura MD (Res), FRCP, FRCPath

Author Information ☺

Journal of Trauma and Acute Care Surgery: December 2021 - Volume 91 - Issue 6 - p 1018-1024
doi: 10.1097/TA.0000000000003360



Original Articles

The Difference in Potential Harms between Whole Blood and Component Blood Transfusion in major Bleeding: A Rapid Systematic Review and Meta-Analysis of RCTs


Louise J Geneen^{a, b, 2, ✉}, Susan J Brunskill^{a, b}, Carolyn Doree^{a, b}, Lise J Estcourt^{a, b, c}, Laura Green^{c, d, e}

Donor selection considerations

We need to mitigate the risk of group O plasma to non-O patients:


- TRALI – male only donors or females tested for HLA/HNA antibodies
 - HTR – select donations that test as low titre for anti-A +B – not everyone is set up to do this routinely
-
- Choice of anticoagulants (CPDA-1 v CPD)
 - D negative or positive, or both?
 - For 'non-standard' LD filters collect in these bags or others?

Donor pool for WB is smaller than for red cells



What are the risks of using D positive blood?

Figure 1. Sequence of events required to cause harm from transfusion of RhD positive red cells in pre-hospital setting




Modelling a change in policy from D neg to D pos for England

Harm	All recipients 1 event every x transfusions (95% CI)	1 child every x transfusions
Major morbidity or mortality due to HTR from index D-positive transfusion	2.7x10 ⁴ (1.1x10 ⁴ – 6.5x10 ⁴)	1.0x10 ⁵ – 9.2x10 ⁴)
Major morbidity or mortality due to future HTR	1.4x10 ⁵ (3.1x10 ⁴ – 3.7 x10 ⁶)	
Foetal death or disability due to HDFN in future pregnancies	1.2x10 ⁴ – 1.2x10 ⁵)	570 (260-2,300)
Any of above	1.4x10 ⁴ (5.6x10 ³ -4.2x10 ⁴)	520 (250-1,700)

<1% improvement in mortality with WB would
 'offset' life years lost due to HDFN risk

1:5 years

Manufacturing considerations

- Time from donation to processing – current IFU for Terumo does not cover 24H ambient hold
 - To LD or not to LD, use of different collection packs & filters
 - Shelf-life
 - The move away from DEHP in the EU
- 
- A thick, solid blue line that curves across the bottom of the slide, starting from the left edge, dipping down towards the center, and then rising towards the right edge.

Special filters are needed to produce D WB containing platelets

Current


Leucodepletion filter
removes platelets



For NHSBT LD WB likely 30% more expensive than RBC + FFP




Plasma



Red
cells


Plasma

1/4 unit of
platelets



Filters approx. 10 x more expensive


Shelf-life is shorter than standard red cells



Shelf life of WB was determined by RBC viability

Historically based on FDA criteria of >75% of RBC remaining in circulation 24 following infusion to healthy volunteers

CPD WB



21 days

CPDA-1 WB



35 days




RBC in AS




35-42 days

Addtiion of adenine and more dextrose increases shelf life either in anticoagulant or AS

It's difficult to know what its shelf life should be




Research Questions

- Is WB superior to component therapy in the resuscitation of bleeding patients?
 - Relevant clinical outcomes – mortality, transfusion requirements
 - Logistical benefits, time on scene
 - What is its optimal shelf-life?
 - Are the risks associated with use of D positive blood acceptable if D negative blood is unavailable?
 - What do patients and the public think?
 - Is it cost-effective?
- 
- A thick, solid blue line that curves horizontally across the bottom of the slide, starting from the left edge, dipping slightly in the middle, and rising towards the right edge.

PPOWER pilot RCT USA n=86

PREHOSPITAL LOW TITER GROUP O WHOLE BLOOD IS FEASIBLE AND SAFE:
RESULTS OF A PROSPECTIVE RANDOMIZED PILOT TRIAL



Injured Patients in
Hemorrhagic Shock

WB vs. Standard Care

Cluster Randomized

Exception From Informed Consent

Feasibility

Halted Enrollment
76% Target

Lower Than Expected
Protocol Adherence
86%

Safety/Efficacy



25% vs 26.1%
28-day Mortality, p=0.8

Lower RBC needs
Lower TEG Abnormalities

TOWAR Study USA

	Prospective RCT
Design	Randomised parallel assignment
Study population	Pre-hospital haemorrhagic shock
Participants	1020
Comparators	Up to 2 units WB pre-hospital versus standard of care
Primary endpoint	30d mortality
Secondary endpoints	Numerous including age of blood (1-14d v >14d)

Recruiting : Due to complete mid 2025

Source: clinicaltrials.gov

TSTORHM study

- France

Design	Prospective RCT non-inferiority
Study population	Trauma triggering MTP
Sample size	164 (82 per arm)
Comparators	WB (up to 6 units) v component therapy
WB used	Terumo filter, 21d shelf-life, 'recycled at 7-14d'
Primary endpoint	MA by TEG
Secondary endpoints	Laboratory markers of haemolysis 2h & 30d mortality Units transfused

SWiFT study

- England

Design	Prospective multi-centre unblinded RCT
Study population	Pre-hospital major traumatic haemorrhage
Sample size	848
Comparators	Wb (up to 2 units) v standard of care (RBC + plasma)
WB used	Terumo filter, 21d shelf-life
Primary endpoint	all-cause mortality or received ≥ 10 units of any blood components ≤ 24 hours from randomisation
Secondary endpoints	Morbidity & Mortality 30, 90 days Hospital resource to discharge Cost-effectiveness Health-related quality of life at 90 days Safety



Thank you!

rebecca.cardigan@nhsbt.nhs.uk

Or connect with me via LinkedIn